

**Quiz**

1. Given a block image  $x_{n_1, n_2}$ , with  $n_1 \in \{0, \dots, N_1 - 1\}$  and  $n_2 \in \{0, \dots, N_2 - 1\}$ , what is the value of  $c_{k_1=0, k_2=0}$ ?
  - A. 1 (whatever the image block is).
  - B. the average of the image block entries, up to a normalizing factor.
  - C. the imaginary part of the image block entries
  - D. the input block turned  $90^\circ$  counter clockwise
2. Consider an image that concatenates  $c_{k_1=0, k_2=0}$  for each block. What does this image look like?
  - A. the same image with a smaller spatial resolution
  - B. the same image with a higher spatial resolution
  - C. the downsampled image
3. Can 2D-DCT decomposition be put into a matrix\*vector form such as  $c = \Phi x$ , where  $\Phi$  is a matrix and  $x, c$  are vectors? and why?

True

False
4. What are the differences/similarities between the classical (sampling+compression) approach and sparse approximation?
5. Which of the following statements are correct?
  - A.  $\Sigma_s$  is a union of subspaces of dimension  $s$
  - B.  $\Sigma_s$  is a union of subspaces of different dimensions
  - C.  $\Sigma_s$  is a subspace of dimension  $s$